

TRB – PG (MATHS)

Complete Study Material

- **Unit Wise - Notes**
- **Unit Wise Questions**
- **Previous Year Question Paper (Problem Solved)**
- **Model Questions**
- **Education (Jeba's Education Guide)**
- **Gk (Special Jeba's GK guide with only full material)**

Unit – VI

Functional Analysis

Banach Spaces - Definition and example - continuous linear transformations - Banach theorem - Natural embedding of X in X^{**} - Open mapping and closed graph theorem - Properties of conjugate of an operator - Hilbert spaces - Orthonormal bases - Conjugate space H - Adjoint of an operator - Projections l^2 as a Hilbert space - l^p space - Holders and Minkowski inequalities - Matrices – Basic operations of matrices - Determinant of a matrix - Determinant and spectrum of an operator - Spectral theorem for operators on a finite dimensional Hilbert space - Regular and singular elements in a Banach Algebra – Topological divisor of zero - Spectrum of an element in a Banach algebra - the formula for the spectral radius radical and semi simplicity.

Banach Spaces

Definition (Normed Linear Space)

- Let N be a linear spaces. A norm on N is a real function $\| \cdot \| : N \rightarrow \mathbb{R}$ satisfying the following conditions:
For $x, y \in N$ and α a scalar,

- $\|x\| \geq 0$ and $\|x\| = 0 \Leftrightarrow x = 0$
- $\|x + y\| \leq \|x\| + \|y\|$
- $\|\alpha x\| = |\alpha| \|x\|$
- A linear space N , with a norm defined on it, is called a Normed linear space.

Definition : Banach Space

- A complete normed linear space is called a Banach space.

Results :

- $\|x\| - \|y\| \leq \|x - y\|$
- Norm is a continuous function on N
i.e., $\|x_n - x\| \rightarrow 0$ as $n \rightarrow \infty$
- In a normed linear space N , addition and scalar multiplication are jointly continuous.
i.e., if $x_n \rightarrow x, y_n \rightarrow y$ and $\alpha_n \rightarrow \alpha$ then
 $x_n + y_n \rightarrow x + y$ and $\alpha_n x_n \rightarrow \alpha x$

Example for Banach Spaces

- The real linear space \mathbb{R} and the complex linear space \mathbb{C} are Banach spaces under the norm defined by $\|x\| = |x|, \forall x \in \mathbb{R} \text{ or } \mathbb{C}$.
- The linear spaces \mathbb{R}^n and \mathbb{C}^n are Banach spaces under the norm defined by
 $\|x\| = (\sum_{i=1}^n |x_i|^2)^{\frac{1}{2}}, \forall x$
 $\mathbb{R}^n \rightarrow n$ – Dimensional Euclidean space
 $\mathbb{C}^n \rightarrow$ Unitary space

3. The space of n-tuples of scalars l_p^n is a Banach space with the norm

$$\|x\|_p = (\sum_{i=1}^n |x_i|^p)^{\frac{1}{p}}$$

4. The linear space l_p of all sequence $x = \{x_1, x_2, \dots, x_n, \dots\}$ of scalars such that $\sum_{n=1}^{\infty} |x_n|^p < \infty$ is a Banach space under the norm defined by

$$\|x\|_p = (\sum_{n=1}^{\infty} |x_n|^p)^{\frac{1}{p}}$$

5. The linear space l_{∞}^n is a Banach space under the norm

$$\|x\|_{\infty} = \max\{|x_1|, |x_2|, \dots, |x_n|\}$$

6. The linear space l_{∞} , of all bounded sequence $x = \{x_1, x_2, \dots, x_n, \dots\}$ of scalar is a Banach space under the norm $\|x\| = \sup_{n=1,2,\dots} |x_n|$

7. The linear space $C(X)$ of all bounded continuous scalar valued functions defined on a topological space is a Banach space under the norm

$$\|f\| = \sup_{x \in X} |f(x)|, f(x) \in C(X)$$

8. The space L_p of all measurable functions defined on a measure space X with the property that $|f(x)|^p$ is integrable is a Banach space under the norm

$$\|f\|_p = \left(\int |f(x)|^p d_m(x) \right)^{\frac{1}{p}}$$

Theorem

Let M be a closed subspace of a normed linear space N and the norm of a coset $x + M$ in the quotient space N/M is defined as $\|x + M\| = \inf\{\|x + m\| : m \in M\}$. Then N/M is a normed linear space. If in addition, if N is a Banach space, then so is N/M .

Holder's Inequality :

$$\sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |y_i|^q \right)^{\frac{1}{q}}$$

$$\text{i.e., } \sum_{i=1}^n |x_i y_i| \leq \|x\|_p \|y\|_q$$

where $p > 1, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$

Minkowski Inequality

$$\left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}}$$

$$\text{i.e., } \|x + y\|_p \leq \|x\|_p + \|y\|_p$$

Definition (Closed unit sphere)

Let X be a Banach space. A closed unit sphere in X is defined as

$$S = \{x \in X : \|x\| \leq 1\}$$

Result : S is a convex set.

Some example for closed sphere:

1. The linear space R^2 , the set of all ordered pairs, is a Banach space under the norm $\|x\| = |x_1| + |x_2|$. The closed unit sphere of R^2 is $S = \{x \in R^2: \|x\| \leq 1\}$ i.e., $S = \{(x_1, x_2) \in R^2: |x_1| + |x_2| \leq 1\}$

2. In R^2 , the norm is defined by

$\|x\|_2 = (|x_1|^2 + |x_2|^2)^{\frac{1}{2}}$ We derive a closed unit sphere.

3. In R^2 , the norm is defined by

$\|x\|_\infty = \max\{|x_1|, |x_2|\}$. We will derive a closed unit sphere.

4. We define the norm

$$\|x\|_p = (|x_1|^p + |x_2|^p)^{\frac{1}{p}}, 1 \leq p < \infty$$

then S is truly spherical iff $p = 2$

In this case, $p < 1$, then

$S = \{x: \|x\| \leq 1\}$ would not be convex.

Definition (Linear transformation)

Let N and N' be two linear spaces over the same system of scalars. A mapping

$T: N \rightarrow N'$ is called a linear transformation if

i) $T(x + y) = T(x) + T(y)$

ii) $T(\alpha x) = \alpha T(x), \forall x, y \in N$

α a scalar

Equivalently,

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y), \forall x, y \in N$$

and α, β scalars

Note : $T(0) = 0$

Definition (Continuous linear transformation)

Let N and N' be two normed linear spaces.

Let $T: N \rightarrow N'$ be a linear transformation if whenever $\{x_n\}$ is a sequence in N such that $x_n \rightarrow x$ in N , then the sequence $\{T(x_n)\}$ converges to $T(x)$ in N'

$$\text{i.e., } x_n \rightarrow x \Rightarrow T(x_n) \rightarrow T(x)$$

Theorem

Let N and N' be two normed linear spaces.

Let $T: N \rightarrow N'$ be a linear transformation, then the following conditions on T are equivalent to one another:

- i) T is continuous
- ii) T is continuous at the origin in the sense that $x_n \rightarrow 0 \Rightarrow T(x_n) \rightarrow 0$
- iii) There exists a real number $K \geq 0$ such that $\|T(x)\| \leq K\|x\|, \forall x \in N$
- iv) If $S = \{x: \|x\| \leq 1\}$ is the closed unit sphere in N , then its image $T(S)$ is a bounded set in N'

Remarks

- i) By condition (iii) of the above theorem, there exists a real number $K \geq 0$ such that

$$\|T(x)\| \leq K\|x\|, \forall x \in N$$

This K is called a bound for T and hence the linear transformation T is bounded linear transformation

- ii) By the above theorem, T is continuous if T is bounded interchangeably

Definition : If T is a continuous linear transformation of N into N' , then norm of T is defined as $\|T\| = \text{Sup}\{\|T(x)\| : \|x\| \leq 1\}$

Remarks

- i) If $N \neq \{0\}$, we can give another equivalent expression of $\|T\|$ as $\|T\| = \text{Sup}\{\|T(x)\| : \|x\| \geq 1\}$
- ii) By the conditions (iii) and (iv) of the above theorem

We arrive at

$$\|T(x)\| \leq \|T\| \|x\|, \forall x \in N$$

Notation

The set of all continuous (bounded) linear transformation from N into N' is denoted by $\mathcal{B}(N, N')$

Theorem

Let N and N' be normed linear space then $\mathcal{B}(N, N')$ is a normed linear space with respect to the pointwise linear operation

$$(T + v)(x) = T(x) + v(x)$$

$$(\alpha T)(x) = \alpha T(x)$$

and with the norm defined by

$$\|T\| = \text{Sup}\{\|T(x)\| : \|x\| \leq 1\}$$

further if N' is a Banach space, then $\mathcal{B}(N, N')$ is also a Banach space.

Definition : Algebra

A linear space N is called an algebra if its element can be multiply in such way that N is a ring and the scalar multiplication is related with the multiplication by the way $\alpha(xy) = (\alpha x)y = x(\alpha y)$

Definition : operator

Let N be a normed linear space. A continuous linear transformation of N into itself is called an operator on N . The set of all operators of a normed linear space N is denoted by $\mathcal{B}(N)$

Results :

1. If N is a Banach space, then $\mathcal{B}(N)$ is also a Banach space
2. $\mathcal{B}(N)$ is an algebra where the multiplication of its elements is defined as $(Tv)(x) = T(vx), \forall v, T \in \mathcal{B}(N)$ and $x \in N$
3. In $\mathcal{B}(N)$, multiplication is jointly continuous i.e.,

$$Tn \rightarrow T \text{ and } Tn' \rightarrow T' \Rightarrow TnTn' \rightarrow TT'$$
4. If $N \neq \{0\}$, then the identity transformation I of N into itself is the identity element of the algebra

$$\mathcal{B}(N) \text{ and } \|I\| = 1$$

Definition

Let N and N' be normed linear spaces

1. An isometric isomorphism of N to N' is a one-to-one linear transformation of N into N' such that $\|T(x)\| = \|x\|, \forall x \in N$
2. N is isometrically isomorphic to N' if T is an isometric isomorphism of N onto N'

Results

1. If M is a closed subspace of a normed linear space N and if T is a natural mapping of N onto N/M defined by $T(x) = x + M, \forall x \in N$, then T is a continuous linear transformation for which $\|T\| \leq 1$
2. If T is a continuous linear transformation from a normed linear space to another normed linear space N' and if M is its null space, then there exists a natural linear transformation T' of N/M into N' which is such that $\|T'\| = \|T\|$

Theorem

Let M be a linear subspace of a normed linear space N and let f be a functional defined on M . If x_0 is not a vector in M and if $M_0 = M + x_0$ is the linear subspace generated by M and x_0 , then f can be extended to a functional f_0 on M_0

such that $\|f\| = \|f_0\|$

Theorem (The Hahn-Banach theorem)

Let M be a linear subspace of a normed linear space N and let f be a functional defined on M . Then f can be extended to a functional f_0 defined on the whole space N such that $\|f\| = \|f_0\|$

Conjugate space of L_p

Let X be a measure space with measure m and p be a general real number such that $1 < p < \infty$. Consider the Banach space of L_p of all measurable functions f defined on X such that $|f(x)|^p$ is integrable

Let g be an element of L_q where $\frac{1}{p} + \frac{1}{q} = 1$

Define a function F_g on L_p by

$$\begin{aligned} F_g(f) &= \int f(x)g(x)d_m(x) \\ |F_g(f)| &= \left| \int f(x)g(x)d_m(x) \right| \\ &\leq \int |f(x)g(x)|d_m(x) \\ |F_g(f)| &\leq \|f\|_p \|g\|_q, \end{aligned}$$

(by Holder's inequality)

Taking Sup for all functions

$f \in L_p$ such that $\|f\|_p \leq 1$

We get $\|F_g\| \leq \|g\|_q$

- It shows that F_g is well defined a scalar valued continuous linear function on L_p with the property that $\|F_g\| \leq \|g\|_q$

- It can be also be shown that every linear functional L_p arises in this way
- Hence $g \rightarrow L_p$ is linear and is an isometric isomorphism of L_q into L_p^*
- This we write as $L_p^* = L_q$

Remarks

1. If we consider that such specialization to the space of all n-tuples of scalars we have $(l_p^n)^* = l_p^*$
If $p = 1$, we have $(L_1^n)^* = L_\infty^n$
2. If we specialize to the space of all sequence of scalar, we have $l_p^* = l_q$
When $p = 1$, $l_1^* = l_\infty$ and $C_0^* = l_1$

Theorem

If N is a normed linear space and if x_0 is a non-zero vector in N , then there exist a functional f_0 in N^* such that

$$f(x_0) = \|x_0\| \text{ and } \|f_0\| = 1$$

Remarks

1. N^* separates vectors in N
i.e., let $x, y \in N$ with $x \neq y$
then there exists a function in N^* such that $f(x) \neq f(y)$

Theorem

If M is a closed linear subspace of a normed linear space N and x_0 is not a vector in M , then there exists a functional f_0 in N^*

$$\text{such that } f(M) = 0 \text{ and } f_0(x_0) \neq 0$$

(i.e., Banach space has rich supply functionals)

Result

Let M be a closed linear subspace of a normed linear space N and let x_0 be a vector not in M . if d is the distance from x_0 to M . Then there exists a functional f_0 in N^* such that

$$f_0(M) = 0 \text{ and } f_0(x_0) = 1 \text{ and } \|f_0\| = \frac{1}{d}$$

Definition (second conjugate space, N^{**})

Let N be a normed linear space. Then the conjugate space of the conjugate space is defined by N^{**} is called the second conjugate space of N

Result

Let N be a normed linear space. Then each vector x in N induces a functional F_x on N^* defined by $F_x(f) = f(x)$ for all $f \in N^*$

$$\text{Such that } \|f_x\| = \|x\|$$

Then the mapping $J: N \rightarrow N^{**}$

(i.e., $x \rightarrow F_x$) defines an isometric isomorphism of N into N^{**}

Remarks

1. The functional F_x , in the above result, is called induced functional.
2. The isometric isomorphism $x \rightarrow F_x$ is called the normal imbedding of N in N^{**} , for it allows us to regard N as a part of N^{**} without altering any of its structure as a normed linear space. Hence we write $N \subseteq N^{**}$

Definition

A normed linear space N is said to be reflexive if $N \equiv N^{**}$

(i.e., the above isometric isomorphism is onto also)

The spaces l_p for $1 < p < \infty$ are reflexive for $l_p^* = l_q$ but $l_q^* = l_p$

$$\text{i.e., } (l_p^*)^* = l_p$$

$$\text{i.e., } l_p^{**} = l_p$$

Remarks

1. Since N^{**} is complete, N is necessarily complete if it is reflexive.
2. But if N is complete, then it need not be reflexive.

For example

$$C_0^* = l_1 \text{ and } C_0^{**} = l_1^* = l_\infty$$

Theorem

If B and B' are Banach space and if T is continuous linear transformation of B onto B' , then the image of each open sphere centred on the origin in B contains an open sphere centred on the origin in B'

Theorem (The open mapping theorem)

If B and B' are Banach space if T is a continuous linear transformation of B onto B' , then T is an open mapping

$$(i.e., G \text{ is open in } B \Rightarrow T(G) \text{ is open in } B')$$

Theorem

A one-to-one continuous linear transformation of one Banach space onto another is a homomorphism.